

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS  
BEFORE OPENING OR USING

  
**WEEDFORCE®**

# Evict® Bio 510

HERBICIDE



**ACTIVE CONSTITUENT: 510 g/L GLYPHOSATE**  
**(present as the isopropylamine salt)**

**GROUP 9 HERBICIDE**

Water soluble herbicide for non-selective control of many annual and perennial weeds in certain situations. Suitable for use in aquatic situations

**IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT**

WEEDFORCE Pty Ltd ACN 602 207 152   
2072 Logan Rd Mt Gravatt QLD 4122  
P: 0488 997 923 [www.weedforce.com.au](http://www.weedforce.com.au)

APVMA Approval No:  
86155/118321

® Trademark of Weed Force Pty Ltd  
© Weed Force Pty Ltd 2017

# WEED FORCE EVICT BIO 510 HERBICIDE

## DIRECTIONS FOR USE

### INDEX

RESTRAINTS.....	4
TABLE 1. PRIOR TO SOWING A CROP OR PASTURE OR TO COMMENCE A FALLOW.....	5
TABLE 2. PASTURE RENOVATION AND TOPPING.....	11
TABLE 3. SUGAR CANE (RATOON CONTROL ) FOR QLD & NSW ONLY .....	12
TABLE 4. RICE -DIRECT DRILLING (NSW ONLY) .....	13
TABLE 5. SORGHUM CONTROL.....	13
TABLE 6. ANNUAL WEEDS TABLE.....	14
TABLE 7. PERENNIAL WEEDS TABLE .....	15
TABLE 8. WOODY WEEDS AND BRUSH TABLE.....	18
AQUATIC WEED CONTROL.....	19
TABLE 9. AQUATIC WEEDS TABLE.....	20
TABLE 10. GENERAL USE SITUATIONS – ALL STATES UNLESS SPECIFIED .....	21
TABLE 11. TANK MIXES WITH METSULFURON (low rate of metsulfuron)	
BY AIR, BOOM, HAND GUN, KNAPSACK.....	23
TABLE 12. TANK MIXES WITH METSULFURON (high rate of metsulfuron)	
BY HAND GUN & KNAPSACK ONLY.....	24
GENERAL INSTRUCTIONS.....	26
CROP ESTABLISHMENT.....	26
MIXING AND APPLICATION.....	26
SURFACTANT.....	28
MIXING.....	28
COMPATABILITY.....	28
EQUIPMENT MAINTENANCE AND USAGE.....	30
RESISTANT WEEDS WARNING.....	30
PROTECTION OF CROP, NATIVE AND OTHER NON-TARGET PLANTS .....	30
PROTECTION OF LIVESTOCK .....	31
PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT .....	31
PRECAUTION .....	31
STORAGE AND DISPOSAL.....	31
SAFETY DIRECTIONS.....	31
FIRST AID .....	31
SAFETY DATA SHEET (SDS).....	31
CONDITIONS OF SALE.....	32

## RESTRAINTS

DO NOT spray if rainfall is expected as rainfall within 6 hours of treatment may reduce the effectiveness of this product. Heavy rainfall within 2 hours of treatment may wash the product from the leaf surface and retreatment may be necessary.

DO NOT disturb treated weeds by grazing, cultivation, sowing etc., after treatment for 1 day for annual weeds and 7 days for perennial weeds to ensure complete uptake of the herbicide; unless specified in the Critical Comments.

DO NOT apply to weeds under stress from frost, cold, waterlogging, lack of moisture. Plants must be actively growing to ensure optimum uptake of the product.

NOTE: Crop and situation, and weeds controlled applies to all states unless stated otherwise.

**TABLE 1. Prior to Sowing a Crop or Pasture or to Commence a Fallow**

Crop and Situation	Weeds Controlled	Rate Vol/ha	Critical Comments
<p>SOUTHERN AUSTRALIA (NSW, VIC, SA, WA, ACT only)</p> <p>Prior to sowing a crop or pasture</p> <p>For weed control prior to sowing a crop or pasture with full soil disturbance by cultivation or sowing with a tyned implement</p>	<p>Barley Grass (<i>Hordeum leporinum</i>), Brome Grass (<i>Bromus unioloides</i>), Volunteer cereals, Wild Oats (<i>Avena</i> spp.)</p>	<p>350 – 700 mL pre tillering</p> <p>700 mL – 900 mL post tillering</p>	<p>Use the higher rate when treating in cold/overcast conditions, when using late in the season. Use the lower rate on young weeds and the higher rate on mature weeds, ie fully tillered grasses or broadleaf weeds at budding or stem elongation.</p> <p>If weeds have been grazed heavily, remove stock prior to spraying to ensure regrowth to 6-8cm before treatment and use the higher rate. Cultivation or planting may proceed from 1 hour of daylight after application to seedling annual weeds if a satisfactory seedbed can be created for crop germination and seedling establishment. If cultivation or sowing does not take place within 21 days, retreatment may be necessary.</p> <p>Annual ryegrass, silver grass and Perennial grasses: A water volume of 70 L/ha or more is recommended with low volume nozzles to improve control. Addition of a non-ionic surfactant according to label directions may improve control.:</p> <p>Crop Establishment: Sowing should not proceed until conditions allow the formation of a satisfactory seedbed. See Crop Establishment for directions.</p> <p>Tank mixtures: for improved control of clovers, add dicamba. Read and follow all label directions for the tank mix products.</p> <p>Perennial Weeds: For Perennial phalaris, Soursob, Skeleton Weed and Sorrel, this product will provide knockdown, seasonal suppression and reduction in treated plant numbers.</p>
	<p>Annual Phalaris (<i>Phalaris canariensis</i>), Annual Ryegrass (<i>Lolium rigidum</i>), Silvergrass (<i>Vulpia</i> spp.) Winter Grass (<i>Poa annua</i>)</p>	<p>700 mL – 900 mL pre tillering</p> <p>900 mL – 1.0L post tillering</p>	
	<p>Capeweed (<i>Arctotheca calendula</i>), Spiny Emex / Doublegee (<i>Emex australis</i>)</p>	<p>350 – 700mL less than 8cm diameter</p> <p>700mL – 900 mL greater than 8cm diameter</p>	
	<p>Amsinckia (<i>Amsinckia</i>), Fumitory (<i>Fumaria officinalis</i>), <i>F. muralis</i>), Paterson's Curse /Salvation Jane (<i>Echium plantaginium</i>), Saffron Thistle (<i>Carthamus lanatus</i>), Scotch Thistle (<i>Onopordum acanthium</i>), Spear Thistle (<i>Cirsium vulgare</i>), Variegated Thistle (<i>Silybum marianum</i>), Volunteer Lupins (<i>Lupinus angustifolius</i>), Wild Turnip (<i>Brassica tournefortii</i>)</p>	<p>700mL – 900 mL less than 12cm diameter</p> <p>900 mL – 1.0L greater than 12cm diameter</p>	
	<p>Dock – seedling (<i>Rumex crispus</i>.)</p>	<p>700 mL – 1.0L</p>	
	<p>Perennial phalaris (<i>Phalaris</i>), Skeleton Weed (<i>Chondrilla juncea</i>) fully emerged rosettes (NSW only)</p> <p>Sorrel (<i>Rumex acetosella</i>), Soursob (<i>Oxalis pes-caprae</i>), Sub Clover (<i>Trifolium subterraneum</i>)</p>	<p>1.0 L</p>	

**TABLE 1.CONTINUED. Prior to Sowing a Crop or Pasture or to Commence a Fallow**

Crop and Situation	Weeds Controlled	Rate Vol/ha	Critical Comments
<p><b>TAS ONLY</b></p> <p>Prior to sowing a crop or pasture</p> <p>For weed control prior to sowing a crop or pasture with full soil disturbance by cultivation or sowing with a tyned implement</p>	All the above weeds	1.0 L – 2.0 L	<p><b>TAS ONLY:</b> Use 1.0 L/ha on annual weeds. Increase to 2.0 L/ha where perennial weeds are being treated. To control White clover and improve control of Sorrel and Dock, add 1 L/ha of Dicamba (200 g/L). Observe dicamba label directions and plant back periods. Addition of wetter at 200 mL/100 L spray solution may improve control.</p>

**TABLE 1.CONTINUED. Prior to Sowing a Crop or Pasture or to Commence a Fallow**

Situation	Weeds Controlled	Rate Vol/ha	Critical Comments
<p><b>SOUTHERN AUSTRALIA</b> (NSW, ACT, SA, Vic, WA only)</p> <p>Before sowing a crop or pasture</p> <p>For weed control prior to sowing a crop or pasture with an implement that gives minimal or no soil disturbance.</p>	Barley Grass ( <i>Hordeum leporinum</i> ), Volunteer cereals, Wild Oats ( <i>Avena</i> spp.)	700 mL – 1.0L	<p>Use the higher rate when treating under cold/ overcast conditions, when using late in the season. Use the lower rate on young weeds and the higher rate on mature weeds, i.e. fully tillered grasses or broadleaf weeds at budding or stem elongation.</p> <p>If weeds have been grazed heavily, remove stock prior to spraying to ensure regrowth to 6-8cm before treatment and use the higher rate.</p> <p><b>Annual Ryegrass, Silvergrass And Perennial Grasses:</b> It is recommended to use a water volume of 70 L/ha or more with low volume nozzles to improve control. Addition of a suitable surfactant at 200 mL/100 L spray solution may improve control.</p>
	Brome Grass ( <i>Bromus unioloides</i> ), Capeweed ( <i>Arctotheca calendula</i> ), Variegated Thistle ( <i>Silybum marianum</i> ), Winter Grass ( <i>Poa annua</i> )	900 mL – 1.4 L	<p>Do not sow if heavy trash is present.</p> <p><b>Aerial Application:</b> May be applied by air if a good seedbed has been established. Always use the higher rates.</p> <p><b>Tank Mixtures:</b> For improved control of Dock, Sorrel and Sub Clover add dicamba. Read and follow all label directions for the tank mix product. Addition of Ammonium Sulphate may improve control when treating under adverse environmental conditions.</p> <p><b>Pasture or Crop Establishment:</b></p> <p>Do NOT sow into excessive trash. Cultivation or planting may proceed from 1 hour of daylight after application to seedling annual weeds if a satisfactory seedbed can be created for crop germination and seedling establishment. Trash may be removed by grazing after treatment. Grazing may commence 6 hours after treatment of annual weeds (small) and 7 days for perennial weeds. Delay grazing for 3 days where annual weeds are large. See also <b>Crop Establishment</b></p> <p><b>Aerial (Or Surface) Seeding:</b> Delay seeding until trash is completely removed by grazing and/ or plant decay. When establishing pasture, ensure application of fertilizer and insecticides and follow-up management is undertaken as required.</p>
	Annual Ryegrass ( <i>Lolium rigidum</i> ), Paterson's Curse /Salvation Jane ( <i>Echium plantaginium</i> ), Saffron Thistle ( <i>Carthamus lanatus</i> ), Scotch Thistle ( <i>Onopordum acanthium</i> ), Silvergrass ( <i>Vulpia</i> spp.), Spear Thistle ( <i>Cirsium vulgare</i> ), Wild Mustard ( <i>Sisymbrium officinale</i> ), Wild Turnip ( <i>Brassica tournefortii</i> )	1.0 L – 1.3 L	
	Erodium ( <i>Erodium cicutarium</i> ), Perennial phalaris ( <i>Phalaris aquatica</i> ), Plantain ( <i>Plantago</i> spp.), Sorrel ( <i>Rumex acetosella</i> ), Sub Clover ( <i>Trifolium subterraneum</i> ),	1.2 L – 1.8 L	
	Dock ( <i>Rumex</i> spp.), Flatweed ( <i>Hyochoeris radicata</i> )	1.8 L	

**TABLE 1.CONTINUED. Prior to Sowing a Crop or Pasture or to Commence a Fallow**

Situation	Weeds Controlled	Rate Vol/ha	Critical Comments
<p><b>TAS ONLY</b></p> <p>Before sowing a crop or pasture</p> <p>For weed control prior to sowing a crop or pasture with an implement that gives minimal or no soil disturbance.</p>	All the above weeds	1.0 – 2.0 L	<p><b>TAS ONLY:</b> Use 1.0 L on annual weeds and increase to 2.0 L on perennial weeds.</p> <p>The product may also be tank mixed with dicamba (1L/ha of 200g/L) to improve control of Sorrel, Dock and White clover. Observe dicamba label directions and plant back periods. Addition of wetter at 200mL/100L spray solution may improve control.</p>

Crop & Situation	Weeds Controlled	Rate Vol/ha	Critical Comments
<p><b>SOUTHERN AUSTRALIA</b> (NSW, ACT, Vic, WA, SA only)</p> <p>For weed control to commence a fallow</p>	<p>Barley Grass (<i>Hordeum leporinum</i>),</p> <p>Volunteer cereals, Wild Oats (<i>Avena</i> spp.)</p>	700 mL – 1.0 L	<p>Use the lower rate on young weeds or where cultivation is to follow within 21 days.</p> <p>Use the higher rate where broadleaf weeds reach stem elongation/budding or where grasses are fully tillered.</p>
	<p>Annual Ryegrass (<i>Lolium rigidum</i>), Brome Grass (<i>Bromus unioloides</i>), Capeweed (<i>Arctotheca calendula</i>), Paterson's Curse /Salvation Jane (rosette) (<i>Echium plantagineum</i>) (rosette), Saffron Thistle (<i>Carthamus lanatus</i>), Scotch Thistle (<i>Onopordum acanthium</i>), Silvergrass (<i>Vulpia</i> spp.), Spear Thistle (<i>Cirsium vulgare</i>), Wild Mustard (<i>Sisymbrium orientale</i>), Wild Radish (<i>Raphanus raphanistrum</i>), Wild Turnip (<i>Brassica tournefortii</i>)</p>	1.0 L - 1.3 L	<p>If weeds have been grazed heavily, remove stock prior to spraying to ensure regrowth to 6-8cm before treatment and use the higher rate.</p> <p><b>Soursob:</b> Treat at tuber exhaustion.</p> <p><b>Hoary Cress:</b> Treat from late rosette to early flowering.</p> <p><b>Annual Ryegrass, Silvergrass and Perennial Grasses:</b> It is recommended to use water volumes of 70L/ha or more with low volume nozzles to improve control.</p> <p>Addition of a suitable surfactant at 200 mL/100 L spray solution wetter may improve control.</p>
	<p>Hoary cress (<i>Cardaria draba</i>),</p> <p>Soursob (<i>Oxalis pes-caprae</i>)</p>	1.0 L	

**TABLE 1.CONTINUED. Prior to Sowing a Crop or Pasture or to Commence a Fallow**

Crop and Situation	Weeds Controlled	Rate Vol/ha	Critical Comments
<p><b>NORTHERN AUSTRALIA</b> (Qld, NSW only)</p> <p>For weed control prior to sowing a summer or winter crop or in a fallow</p>	<p>Annual Phalaris (<i>Phalaris canariensis</i>), Barley Grass (<i>Hordeum leporinum</i>), Volunteer cereals, Wild Oats (<i>Avena</i> spp.)</p>	350 - 700mL	<p>Use the lower rate on young weeds or where cultivation is to take place within 21 days.</p> <p>Use the higher rate where broadleaf weeds reach stem elongation/budding or where grasses are fully tillered.</p>
	<p>Barnyard Grass (<i>Echinochloa crus-galli</i>), Liverseed Grass (<i>Urochloa</i> spp.), Stinkgrass/ Lovegrass (<i>Eragrostis curvula</i>)</p> <p>Sweet Summer Grass, Volunteer sorghum (<i>Sorghum halapense</i>)</p>	700mL –1.4 L	<p>At more advanced stages certain broadleaf weeds may require the higher rate range or the addition of 2,4-D.</p> <p>In winter (cold) conditions, symptoms on Deadnettle may be slow to develop.</p> <p>If weeds have been grazed heavily remove stock prior to spraying to ensure regrowth to 6-8cm before treatment and use higher rate.</p>
	<p>Australian bluebell (Qld only) (<i>Wahlenbergia gracilis</i>), Cudweed (<i>Gnappalium luteoalbum</i>),Fumitory (<i>Fumaria officinalis</i>, <i>F. muralis</i>), Mexican poppy (<i>Argemone ochroleuca</i>), Mintweed, (<i>Salvia reflexa</i>)</p> <p>New Zealand spinach (<i>Tetragonia tetragonoides</i>),</p> <p>*Noogoora Burr (<i>Xanthium pungens</i>) Saffron Thistle (<i>Carthamus lanatus</i>), Spear Thistle (<i>Cirsium vulgare</i>), Spurge <i>Euphorbia</i> spp.),</p> <p>* Variegated Thistle (<i>Silybium marianum</i>)</p> <p>*Volunteer sunflower, Yellowwine/Caltrop (<i>Tribulus terrestris</i>)</p> <p>Wireweed (<i>Polygonum aviculare</i>)</p>	700 mL - 1.0 L	<p>Liverseed and Barnyard grass may be sensitive to moisture stress. Dense stands may require re-treatment.</p> <p><b>Aerial Application:</b> See General Instructions. Do not apply by air if temperature is over 30°C.</p> <p>*Larger plants (&gt;5cm) of Noogoora Burr, Variegated Thistle and Volunteer Sunflower may require up to 1.3 L/ha to achieve control.</p> <p><b>Crop Establishment:</b> Sowing should not proceed until conditions allow for a formation of a satisfactory seedbed.</p> <p>See Crop Establishment for directions.</p> <p><b>Sowthistle:</b> Previously grazed plants may be difficult to control without allowing full recovery.</p>

**TABLE 1.CONTINUED. Prior to Sowing a Crop or Pasture or to Commence a Fallow**

Crop and Situation	Weeds Controlled	Rate Vol/ha	Critical Comments
<b>NORTHERN AUSTRALIA</b> (Qld, NSW only)  For weed control prior to sowing a summer or winter crop or in a fallow (cont)	Boggabri weed ( <i>Amaranthus macrocarpus</i> ), Caltrop ( <i>Tribulis terrestris</i> ), Indian hedge mustard ( <i>Sisymbrium orientale</i> ), Mintweed ( <i>Salvia reflexa</i> ), Summer grass ( <i>Digitaria ciliaris</i> )	350 – 700 mL up to 5 true leaves or 3cm in diameter or height  or  700 mL – 1.0 L if greater than 3cm in diameter or height	Continued from above
	Annual ground cherry ( <i>Physalis ixocarpa</i> ), Bladder Ketmia, Sow Thistle ( <i>Sonchus oleraceus</i> )  Turnip weed ( <i>Rapistrum rugosum</i> ),  Wild Lettuce ( <i>Lactuca saligna</i> )  Wild Turnip ( <i>Brassica tournefortii</i> )	700 mL – 1.0 L prior to stem elongation/ budding  Or 1.0 – 1.3 L after stem elongation/ budding	

**TABLE 2. Pasture Renovation and Topping**

Crop and Situation	Weeds Controlled	Rate (L/ha)	Critical Comments
<b>Pasture with Poa Tussock present as a weed.</b> (QLD, NSW, ACT, Tas, VIC, only)  For pasture renovation	Annual weeds (see previous table) and Poa tussock ( <i>Poa labillardieri</i> )	2.1 – 2.8 L	<b>Before spraying:</b> * Graze heavily * Remove stock 14 days or more before treatment * Apply to actively growing plants after the autumn break but before heavy frosts (March – May)  Increasing to the higher rate may give more effective reductions.  Sowing of new pasture may begin 14 days after treatment. It is essential that correct follow-up pasture establishment and management occur after treatment. Spot treatment will limit re-infestations. May be aerially applied.
<b>Pasture with Bent Grass present as a weed.</b> (Vic, Tas only)  For control/ suppression of Bent Grass prior to sowing a crop or pasture.	Annual weeds (see previous table) and Bent grass ( <i>Agrostis tenuis</i> )	1.8 L	Apply late spring when seed-heads have development, but before the onset of summer moisture stress.  Remove stock prior to spraying to achieve good foliage coverage. Ensure plants are actively growing.  <b>FOLLOW-UP MANAGEMENT:</b> 10-21 days after spraying, fully disturb soil with a tined implement and then sow summer crop and/or reseed pasture or crop the following autumn.
<b>Pasture Topping</b> (WA, SA, Vic, Tas, NSW, ACT, only)  for the reduction of seed set of annual grass, Capeweed and Calomba daisy	Annual Ryegrass ( <i>Lolium rigidum</i> ), Calomba Daisy ( <i>Pentzia sufruticosa</i> )	330 mL	Use the higher rate for dense infestations or where Annual Ryegrass is present. Apply before signs of plants 'haying off' .
	Barley Grass ( <i>Hordeum leporinum</i> ), Brome Grass ( <i>Bromus unioloides</i> ), Capeweed ( <i>Arctotheca calendula</i> ), Silvergrass ( <i>Vulpia</i> spp.)	210 – 330 mL	Annual Ryegrass and Capeweed - Apply at flowering.  Other weeds – apply from head to milky dough stage.  Stock should be removed before spraying to allow regrowth.  Pasture legumes may be affected. DO NOT apply to clover or medic/clover crops to be used for seed or hay. Apply a maximum of 50 L/ha water. Above this water volume add a non-ionic surfactant

**TABLE 2 CONTINUED. Pasture Renovation and Topping**

Crop and Situation	Weeds Controlled	Rate (L/ha)	Critical Comments
<b>Pasture Manipulation</b> (WA, NSW, VIC, ACT only)  for control / suppression of certain pasture grasses before sowing soybeans, forage crops or Leucaena.	Carpet grass ( <i>Axonopus</i> spp.), Kikuyu ( <i>Pennisetum clandestinum</i> ), Paspalum ( <i>Paspalum dilatatum</i> )	1.0 – 4.2 L	Apply the lower rate for the suppression only. The higher rate will provide control.  <b>Leucaena</b> (Qld ONLY):  Rows should be 4m apart. Use 1.8 L/ha with asingle taper fan nozzle LF1-80 mounted at the rear of the single row planter giving a 1m swath.
	<b>Pasture Manipulation</b> (QLD only)  for control / suppression of certain pasture grasses before sowing soybeans, forage crops or Leucaena.	Carpet grass, ( <i>Axonopus</i> spp.) Paspalum ( <i>Paspalum dilatatum</i> )	
	Kikuyu ( <i>Pennisetum clandestinum</i> )	400 mL – 4.2 L	
	Black Speargrass ( <i>Hederopogon confortus</i> ), Wire grasses ( <i>Anstida</i> spp.), Love grasses ( <i>Eragrostis</i> spp.), Red Natal grass ( <i>Rhynchelytrum repens</i> )  Barbed wire grass ( <i>Cymbopogon refractus</i> )	2.1 L	

**TABLE 3. Sugar Cane (Ratoon Control ) For QLD & NSW Only**

Crop and Situation	Variety	Rate (L/ha)	Critical Comments
<b>Sugar Cane</b>  Ratoon control	Q63, Q87, Q90, Q117, Q120, Q129, Q130, H56-752, Pindar, Triton	2.1 – 2.8 L	Apply when ratoons are actively growing and are 60 – 100cm tall. Use the lower rate for suppression or where control by cultivation is planned. Use the higher rate for control.
	Q86, Q96, Q113	2.8 – 3.5 L	
	Cassius, Q115, Q122, Q94	3.5 – 4.2 L	Apply with properly calibrated boom sprayer at correct height to allow overlap of the spray pattern at the top of the crop canopy.  DO NOT apply if plants are under stress from waterlogging or low moisture.
	NC0310, Q107	4.2 – 6.3 L	

**TABLE 4. Rice – Direct Drilling for NSW Only**

Crop and Situation	Weeds Controlled	Rate (L/ha)	Critical Comments
<b>Rice</b>  Direct Drilling	Annual Ryegrass ( <i>Lolium rigidum</i> ), Annual Phalaris ( <i>Phalaris canariensis</i> ), Barley Grass ( <i>Hordeum leporinum</i> ), Burr Medic ( <i>Medicago</i> spp.), Sub Clover ( <i>Trifolium subterraneum</i> ), Winter Grass ( <i>Poa annua</i> )	700 mL – 900 mL	If plants are drought stressed a pre-watering must be applied. If the site has been grazed, allow plants to regrow to 6-8cm before treatment.  For the control of Annual Ryegrass use the higher rate and add a suitable non-ionic wetting agent at the recommended rate.  <b>Crop Sowing</b> – Sow 1 – 14 days after treatment.  Residual control will only be achieved by adding another suitable herbicide.

**TABLE 5. Sorghum Control**

Crop and Situation	Weeds Controlled	Rate (L/ha)	Critical Comments
<b>Sorghum Control</b> (Qld, NSW, only)  Before harvest	Grain sorghum ( <i>Sorghum bicolor</i> )	1.0 or 1.3L	DO NOT apply to varieties intended for seed production or varieties prone to lodging.  DO NOT apply to crops under stress from factors such as waterlogging, frost, disease, low moisture etc.  Apply when grain moisture is less than 25%. The product can be applied when some browning has occurred.  Use the lower rate for control of the crop, late tillers and suppression of ratoon regrowth. Use the higher rate for better suppression of ratoon regrowth.  Treatment may increase potential for crop lodging, particularly if prior moisture stress has occurred.  Harvest as soon as sufficient dry-down has occurred to prevent further lodging.  CAUTION: Sorghum may be naturally toxic to stock.
<b>Sorghum Control</b> (Qld, NSW, only)  Post-harvest	Sorghum stubble (Grain sorghum) ( <i>Sorghum bicolor</i> )	700 mL – 1.0L for new regrowth from slashed stubble  1.2 – 1.6 L for standing green stubble  800mL – 1.2 L for fresh spring growth	DO NOT apply to crop under stress from such factors as waterlogging, frost, disease, low moisture etc. For slashed stubble and spring regrowth apply when regrowth is at least 20cm high.  Standing Stubble - apply only if sufficient green leaf is present. Allow regrowth of at least to 20cm if grazing has occurred.  Use the lower rate for knockdown and regrowth suppression where cultivation is to follow.  Use the higher rate for better control of regrowth. It is important to note that variable results can occur if the crop has been under stress or grown under marginal conditions. The varieties Ruby, Trump, Nugget2, Goldrush 2 and Prize are particularly susceptible if growing conditions are not ideal.  CAUTION: Sorghum may be naturally toxic to stock.

**TABLE 6. Annual Weeds – For All States**

Weeds Controlled	Rate	Critical comments
Amaranth ( <i>Amaranthus</i> spp.) Barley Grass ( <i>Hordeum leporinum</i> ) Barnyard Grass ( <i>Echinochloa crus-galli</i> ) Brome Grass ( <i>Bromus</i> spp.) Caltrop ( <i>Tribulus terrestris</i> ) Canary Grass ( <i>Phalaris</i> spp.) Capeweed ( <i>Arctotheca calendula</i> ) Cereals Volunteer Chickweed ( <i>Stellaria media</i> ) Cobbler's Peg ( <i>Bidens pilosa</i> ) Fumitory ( <i>Fumaria officinalis</i> ) Ground Cherry ( <i>Physalis angulata</i> ) Lesser Swinecress ( <i>Coronopus didymus</i> ) Liverseed Grass ( <i>Urochloa panicoides</i> ) Mintweed ( <i>Salvia reflexa</i> ) Paradoxa grass ( <i>Phalaris paradoxa</i> ) Paterson's Curse ( <i>Echium plantagineum</i> ) Pigweed ( <i>Portulaca oleracea</i> ) Potato weed ( <i>Galinisoga parviflora</i> ) Ryegrass Annual ( <i>Lolium rigidum</i> ) Saffron Thistle ( <i>Carthamus lanatus</i> ) Silver grass ( <i>Vulpia</i> spp.) Sow Thistle ( <i>Sonchus oleraceus</i> ) Spear Thistle ( <i>Cirsium vulgare</i> ) Spiny Burrgrass ( <i>Cenchrus</i> spp.) Spurge ( <i>Euphorbia</i> spp.) Sub Clover ( <i>Trifolium subterraneum</i> ) Wild Mustard ( <i>Sisymbrium officinale</i> ) Wild Oats ( <i>Avena</i> spp.) Wild Turnip ( <i>Brassica tournefortii</i> ) Winter Grass ( <i>Poa annua</i> ) Variegated Thistle ( <i>Silybum marianum</i> )	<p><b>Boom:</b> 1.3 – 2.1 L/ha</p> <p><b>Handgun:</b> 350 – 490mL per 100L of water</p> <p><b>Knapsack:</b> 50 – 70mL per 15L of water</p>	<ul style="list-style-type: none"> <li>Apply only to plants which are actively growing and not suffering stress.</li> <li>Use the lower rate for weeds up to 15cm tall; and the higher rate for weeds over 15cm tall.</li> <li>The effects of the product may take 3-7 days to appear under normal conditions and up to 20-30 days in cool conditions.</li> <li>No residual control will be provided by this product. Germinations after initial treatment may have to be re-sprayed.</li> </ul> <p>For residual control, this product should be tank mixed with a suitable residual herbicide.</p>

**TABLE 7. Perennial Weeds**

Weeds Controlled	Rate			Critical Comments
	Boom L/ha	Knapsack mL/15L	Handgun vol/100L	
Bamboo ( <i>Bambusa</i> spp.)	-	100 mL	700 mL	Apply to actively growing foliage and/or regrowth which is between 1m and 2m tall.  <b>Cut Stump:</b> Dilute 1:6 i.e. mix 1 part of this product with 6 parts water. Cut stems back to 20cm high, pour mixture down hollow stem or wet the cut.
Bent grass (Vic, Tas only) <i>(Agrostis tenuis)</i>	1.8 L	50 mL	350 mL	Apply to plants in late spring which have some seed head development late in the spring. Plants must be actively growing. It is necessary to follow-up spraying with full soil disturbance within 21 days and then plant to a summer crop and/or by reseeding pasture or crop in autumn.
Blady grass (NSW, ACT, Qld only) <i>(Imperata cylindrica)</i>	6.3 L	140 mL	900 mL	Apply to actively growing plants when most plants have reached head stage.
Bracken <i>(Pteridium esculentum)</i>	-	160 mL	1 L	<ul style="list-style-type: none"> <li>For best control, Wiper application is recommended.</li> <li>Bracken should be slashed in the previous winter/spring so that application is made to new growth.</li> <li>Apply to actively growing, fully unfurled fronds in autumn (March-May) before the onset of frosts.</li> <li>Symptoms may be slow to appear.</li> <li>Follow-up treatment is recommended as control will NOT be achieved after one treatment.</li> </ul>
Carpet grass <i>(Axonopus</i> spp)	2.1 L	50 mL	350 mL	Apply to actively growing plants at early-head stage.
Cocksfoot <i>(Dactylis glomerata)</i>	2.1 L	70 mL	500 mL	Apply to actively growing plants at early-head stage.
Couch <i>(Cynodon dactylon)</i>	6.3 L	130 mL	900 mL	Apply to actively growing plants when most are at the early-head stage. For best results in WA and SA apply in October-November.
Flatweed (Cat's Ear) <i>(Hypochaeris radicata)</i>	2.1 L	70 mL	500 mL	Apply at early flowering to fully developed rosettes.
Guinea grass <i>(Panicum maximum)</i>	6.3 L	130 mL	900 mL	Apply to actively growing plants at early head stage. May be applied by 'Wiper Equipment'



TABLE 7 CONTINUED. Perennial Weeds

Weeds Controlled	Rate			Critical Comments
	Boom L/ha	Knapsack mL/15L	Handgun vol/100L	
Hoary Cress (NSW, ACT, Tas, Vic only) <i>(Cardaria draba)</i>	1.0 L	50 mL	350 mL	Apply late July to September to actively growing plants at late rosette to flowering stage.  Ensure plants are not stressed at time of spraying. Where stems are long enough, Wiper Equipment may be used.  Tas: add a non-ionic surfactant at the recommended rate.
Johnson grass <i>(Sorghum halepense)</i> Kangaroo grass <i>(Themeda australis)</i> Kikuyu grass <i>(Pennisetum clandestinum)</i>	4.2 L	100 mL	700 mL	Apply to actively growing plants at early-head stage. May be applied by Wiper Equipment to Johnson Grass.
Lovegrass, African (Vic, NSW, ACT, WA only) <i>(Eragrostis curvula)</i>	4.2 L	100 mL	700 mL	Apply to actively growing plants. To restrict seedling re-establishment, pasture improvement is recommended
Nutgrass <i>(Cyperus rotundus)</i>	4.2 L	100 mL	700 mL	<b>Non-cultivated situations.</b> Apply to actively growing plants in February -April.
Nutgrass <i>(Cyperus rotundus)</i>	2.1 L followed by 2.1 L	70 mL followed by 70 mL	500 mL followed by 500 mL	<b>Cultivated situations:</b> Make first application when at least 20% of plants have reached early-head stage (about Feb). Make the second application when most plants have re-emerged (about 6-8 weeks after the first application). Follow-up treatments may be necessary as further plants emerge.
Pampas grass <i>(Cortaderia spp)</i>	-	100 mL or 130 mL	700 mL or 900 mL	Apply to actively growing plants during spring, summer or autumn. Ensure complete coverage of the foliage. For best results apply at flowering. Use the lower rate for plants under 1m tall and the higher rate for larger plants. Plants may be cut prior to application but regrowth must be at least 1m prior to spraying.
Paragrass <i>(Brachiara mulica)</i>	6.3 L	130 mL	900mL	Apply to actively growing plants at the early head stage.
Paspalum <i>(Paspalum dilatatum)</i>	4.2 L	100 mL	700 mL	Apply to actively growing plants at the early head stage.
Phalaris (SA, Vic, NSW, ACT only) <i>(Phalaris aquatica)</i>	2.1 L or 4.2 L	50 mL or 100 mL	350 mL or 700mL	Apply in winter/spring to actively growing plants. Use lower rate when only knockdown is required such as prior to burning for a firebreak. Burning should not take place for 2-3 weeks after spraying. The higher rate should be used for longer-term control.

TABLE 7 CONTINUED. Perennial Weeds

Weeds Controlled	Rate			Critical Comments
	Boom L/ha	Knapsack mL/15L	Handgun vol/100L	
Plantains <i>(Plantago spp)</i>	2.1 L	70 mL	500 mL	Apply to actively growing plants at early head stage. Symptoms may be slow to appear
Prairie grass <i>(Bromus unioloides)</i> Old Blue grass <i>(Dichanthium sericium)</i> Redleg grass <i>(Bothriochloa ambigua)</i> Rhodes grass <i>(Chloris gayana)</i>	4.2 L	100 mL	700 mL	Apply to actively growing plants at early head stage
Rope Twitch (Tas, Vic only) <i>(Agropyron repens)</i>	4.2 L	100 mL	700 mL	Apply in late summer-autumn to actively growing plants with foliage at least 20cm high. To ensure maximum root emergence the area should not be cultivated in the period from the preceding winter until the time of spraying.
Sorrel <i>(Rumex acetosella)</i>	4.2 L	100 mL	700 mL	Apply to actively growing plants when the majority of the plants are at the early bud stage
Soursob (NSW, ACT, Vic, SA, WA Tas only) <i>(Oxalis pes-caprae)</i>	1.0 L	50 mL	350 mL	Apply to actively growing plants late July to early September prior to plant senescence (yellowing). Ensure plants are not stressed at time of application. If plants have been grazed or frosted, allow regrowth before treatment.
St John's Wort <i>(Hypericum perforatum)</i>	2.1 L	50 mL	350 mL	Apply to actively growing plants at flowering to post-flowering, procumbent stem stage (about Nov-May). Pasture improvement or re-treatment may be necessary to prevent seedling re-establishment.
Thistle Artichoke (Vic, SA only) <i>(Cynara cardunculus)</i>	2.1 L	50 mL	350 mL	Apply when plants are at the rosette to early head stage.
Thistle Californian (Vic, Tas only) <i>(Cirsium arvense)</i>	4.2 L	50 mL	350 mL	Apply to actively growing plants at the flowering stage. To ensure maximum shoot emergence the area should not be cultivated prior to spraying. Re-treatment and/ or pasture improvement may be necessary to restrict seedling re-establishment.
Yorkshire fog <i>(Holcus lanatus)</i>	2.1 L	70 mL	500 mL	Apply to actively growing plants

**TABLE 8. Woody Weeds and Brush**

Weeds Controlled	Rate		Critical Comments <b>Read Application Checklist before using</b>
	Handgun, mL/100L	Knapsack, mL/15L	
Bitou Bush/ Boneseed (NSW, ACT, Qld, Vic, Tas only) ( <i>Chrysanthemoides monilifera</i> )	350 or 700 mL	50 or 100 mL	Apply to actively growing plants. Do not treat plants that are stressed, particularly drought stressed. Spray to wet all foliage. Best results are achieved when treated during the winter at peak flowering.  Use the higher rate on larger bushes. Follow-up treatment may be required to prevent the establishment of germinating weeds.
Blackberry ( <i>Rubus fruticosus</i> )	700 or 900mL	100 or 130 mL	Apply from January to May (flowering to leaf fall). Spray plants which are not under stress to thoroughly wet all foliage. Use the higher rate for dense old stands over 2m high. Further treatment may be needed to control seedlings and regrowth. Symptoms may be slow to appear and may not be apparent until the next season. TAS ONLY: Do not spray bushes bearing mature fruit.
Boxthorn ( <i>Lycium ferocissium</i> )	500 or 700 mLmL	70 or 100 mL	Spray to wet all foliage. Use the lower rate for young bushes and the higher rate for bigger mature bushes. DO NOT spray if conditions are hot and dry. Regrowth and seedling germination may have to be retreated.
Crofton Weed (NSW, ACT, Qld only) ( <i>Eupatorium adenophorum</i> )	350 mL	50 mL	Apply to plants with full foliage, which are actively growing. Spray to wet all foliage. Seedling germination may have to be retreated.
Groundsel Bush (NSW, ACT, Qld only) ( <i>Baccharis halimifolia</i> )	500 or 700 mL	70 or 100 mL	Apply to actively growing plants using the higher rate for plants over 2m tall. Do not spray during summer drought conditions or in winter. Spray to wet all foliage. Seedling germination may have to be retreated.
Hawthorn (NSW, ACT, Vic, Tas, WA, SA only) ( <i>Cratagus</i> spp.)	700 or 900 mL	100 or 130 mL	Spray from flowering to leaf fall when plants are actively growing. Use the higher rate on plants over 2m tall. Spray to thoroughly wet all foliage. Seedling regrowth may have to be retreated
Lantana (NSW, ACT, Qld only) ( <i>Lantana camara</i> )	700 mL	100 mL	Apply to plants with full foliage, which are actively growing. Spray to thoroughly wet all foliage and individual plants. Seedling regrowth may have to be retreated
Mistflower (NSW, ACT, Qld only) ( <i>Eupatorium riparium</i> )	350 mL	50 mL	Apply to plants with full foliage, which are actively growing. Spray to thoroughly wet all foliage. Seedling regrowth may have to be retreated
Sifton Bush/ Chinese Scrub (NSW, ACT, Qld only) ( <i>Cassubua arcutata</i> )	700 or 900 mL	100 or 130 mL	Apply to actively growing plants ensuring complete coverage. Seedling regrowth may have to be retreated. For high-volume application, use the higher rate when bushes are over 1m.  For Wiper Application a double pass application is required. Best results are achieved if bushes are less than 1m tall and green at the time of application.
Sweet Briar (NSW, ACT, Vic, Tas, WA, SA only) ( <i>Rosa rubiginosa</i> )	1.0L or 1.3L	160 or 210 mL	Apply from late flowering to leaf fall to actively growing plants. Spray to thoroughly wet all foliage. Use the higher rate for bushes over 1.5m tall. Seedling regrowth may have to be retreated.

**AQUATIC WEED CONTROL**

Reduction in effectiveness may result if more than ¼ of the aboveground portion of the weed is submerged at treatment. Submerging the treated plants following treatment may result in the spray being washed from the plant surface, thus reducing effectiveness. DO NOT apply this product within 0.5km of potable water intake in flowing water (e.g. river or stream) or within 0.5km of a potable water intake in a standing body of water such as lake, pond or reservoir. Applications to moving bodies of water should be made while travelling upstream wherever possible to prevent concentration of this herbicide in water. When making bankside applications, do not overspray more than 0.5m into open water. Avoid spraying across moving bodies of water where weeds do not exist.

**DO NOT ADD EXTRA SURFACTANT/WETTER, UNLESS IT IS APPROVED IN AQUATIC SITUATIONS.**

When spraying floating weeds, use a low volume, low pressure boom sprayer or sprinkler sprayer. DO NOT submerge weeds when spraying as this may wash herbicide off the leaves. When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid sudden impact on the habitat.

**TABLE 9. Aquatic Weeds**

Weeds Controlled	Rate			Critical Comments
	Boom L/ha	Knapsack mL/15L	Handgun vol/100L	
Alligator weed	4.2 L	110 mL	700 mL	Apply to actively growing plants from Summer through Winter. Floating form only.
Brown Beetle Grass (NSW, ACT only)	2.1 L	55 mL	350 mL	Apply to actively growing plants. Do not apply to partially submerged plants.
Cumbungi ( <i>Typha</i> spp.)	6.3 L	140mL	900 mL	Spray during Summer or Autumn period during the heading stage. Except for Tasmania, Wiper equipment can be used. Refer to information on 'Application Equipment' section of the label.
Paragrass ( <i>Bracharia mutica</i> )	6.3 L	140 mL	900mL	Spray at early head stage when plants are in active growth.
Phragmites, Common Reed ( <i>Phragmites australis</i> )	6.3 L	140 mL	900 mL	If the Wiper technique is to be used, refer to 'Wiper Equipment' section of this booklet. Spray when plants are getting close to early head stage and actively growing. Spray symptoms may not be observed for a season or more.
Rushes ( <i>Juncus</i> spp)	See Critical Comments			Use Wiper technique ensuring a high percentage of green matter is present. Refer to section of this booklet entitled 'Wiper Equipment' for directions for use.
Sedge – Tall (NSW, ACT, Tas, Vic only) ( <i>Cyperus gracilis</i> )				
Water Couch ( <i>Paspalum distichum</i> )	6.3 L	140 mL	900mL	Spray actively growing plants in February/March period. 75% of plants should be visible above the water at the time of spraying.
Water Hyacinth ( <i>Eichornia crassipes</i> )	4.2 to 6.3 L	110 mL to 140 mL	700 mL to 900mL	Spray actively growing plants at or beyond the early bloom stage of growth. Use the higher rate on dense infestations.
Water Lettuce ( <i>Pistia stratiotes</i> )	-			Best results are obtained from mid-Summer through to Winter. Use the higher rate on dense infestations.
Waterlily, Yellow ( <i>Nymphaea mexicana</i> )	4.2 L	110 mL	-	Apply when there is a maximum emergence of floating leaves. Allow 2-3 weeks for symptoms to develop, and then retreat any unaffected plants. Use low volume sprayer.

**TABLE 10. General Use Situations – all states unless specified.**

Situation	Weeds Controlled	Application Rates	Critical Comments
<b>Agricultural Areas</b>	See Weeds Controlled table for Annual and Perennial Weeds, and Brush and Woody weeds.	See Weeds Controlled Tables	For the control of weeds listed in "Weeds Controlled" prior to sowing of crop.
		7 mL per litre of water	Ensure weeds are actively growing at time of application. Complete and uniform coverage is necessary to ensure best results. Symptoms may take 3-21 days to appear. NO residual control is provided.
<b>Domestic areas, Commercial, Industrial areas, Public service areas, Rights-of-way,</b>			
<b>Forestry Situations</b>		See Weeds Controlled Tables	This product may be used: <ol style="list-style-type: none"> <li>In site preparation before planting.</li> <li>Before establishment of nurseries.</li> <li>Amongst established trees by using selective wiper equipment, directed or shielded spray. This product must not contact foliage or green bark of desirable trees. The wiper should not contact any part of the tree.</li> </ol>
<b>Pastures</b>			This product may be used by the following methods: <b>Spot Application</b> – to remove weeds by spot application within a pasture. This product is non-selective and may damage or kill any plant in the sprayed area. To prevent seedling establishment, pasture improvement and/or re-treatment may be necessary. <b>Boom Application</b> – This product may be used to suppress or kill existing pasture prior to reseeding or establishment of other crops. <b>Selective Application</b> - See Wiper Equipment under General Instructions.)

**TABLE 10 CONTINUED. General Use Situations – all states unless specified.**

Situation	Weeds Controlled	Application Rates	Critical Comments
<p><b>Horticultural Crops</b></p> <p>Avocado, Banana, Blueberries, Citrus fruit, Custard apples, Duboisia Figs – dessert, Guava, Kiwifruit, Litchi, Mango, Monstera fruit, Nuts (including Almond, Pecan, Macadamia, Pistachio and Walnut).</p> <p>Olives, Pawpaw, Persimmons, Pome fruit, Raspberries, Stone fruit, Tea, Vineyards.</p>	See Weeds Controlled table for Annual and Perennial Weeds, and Brush and Woody weeds.	See Weeds Controlled Tables	<p>This product can be used as a shielded or directed spray, or using wiper equipment. DO NOT apply near trees or vines less than 3 years old unless they are adequately protected from spray and spray drift. DO NOT allow spray or spray drift to contact bark, leaves, wounds or any other plant parts of any crop as severe injury may occur.</p> <p><b>Tea:</b> Apply a maximum of 2.8 L/ha by shielded spray or directed off-centre nozzle or 350 mL/100L by directed handgun or knapsack to avoid injury to the crop.</p>
<p>Peanuts, Cotton, Soybeans, Sugar Cane</p> <p>Using selective application equipment only.</p> <p>QLD, NSW only</p>			<p><b>WIPER EQUIPMENT</b> Apply to the weeds growing between the rows or to weeds growing at least 15cm above the crop. DO NOT allow the product to contact the crop or to drip from the applicator as serious crop injury may occur.</p> <p><b>SHIELDED SPRAYERS</b> (Cotton only) Apply to the weeds growing between the rows using a shielded sprayer. DO NOT apply unless the crop is at least 20cm high. Do not allow product or drift to contact crop.</p>

**NATIVE PASTURES, RIGHTS OF WAY, COMMERCIAL AND INDUSTRIAL AREAS, DOMESTIC AND PUBLIC SERVICE AREAS**

**CONTROL USING HIGHER RATE OF GLYPHOSATE AND LOWER RATE OF METSULFURON**

**- WEED FORCE EVICT Bio 510 + RENOVATE BROADLEAF & WOODY WEED TANK MIX (metsulfuron-methyl 600g/kg)**

**(always refer to First Aid, Safety Directions, Resistant Weeds Warning and any other restrictions for Weed Force Renovate Herbicide)**

**FOR ALL STATES**

**RESTRAINTS:**

**DO NOT** apply if rainfall is expected within 6 hours of spraying.

**DO NOT** store tank mixes with other products.

**DO NOT** store the prepared spray for more than 2 days.

**TABLE 11. Control using higher rate of glyphosate and lower rate of metsulfuron**

Situation	Weeds Controlled	Rate ALWAYS ADD HOLDFAST ORGANOSILICONE PENETRANT		Critical Comments <b>For All Weeds:</b> Apply when target weed is in active growth and not under stress from water logging, drought etc. Results may be slow to appear and may not be visible for 3-6 weeks. Full brown-out may take 3-6 months.
		Boom or Aerial	Hand Gun or Knapsack	
Native Pastures **, Forestry, Rights of Way, Commercial and Industrial Areas, Domestic and Public Service Areas	Blackberry ( <i>Rubus spp</i> )	5.6L + 60 g Renovate per ha	280 mL + 3 g Renovate per 100 L water	For Blackberries, apply from flowering until prior to leaf yellowing. Due to widespread picking of Blackberries by the public, it is not recommended that the product be applied to bushes bearing mature fruit. Use Holdfast Organosilicone penetrant at the rate of 200-500 mL per 100 L of water
	Bracken ( <i>Pteridium esculentum</i> )	2.8 L + 30 g Renovate per ha		For Bracken, apply when fronds are fully unfurled but prior to first frosts. For boom application, refer to boom application section. Use Holdfast Organosilicone penetrant at the rate of 200-500 mL per 100 L of water
	Gorse ( <i>Ulex europaeus</i> )			For Gorse, apply when actively growing at any time of the year, except Spring. Use Holdfast Organosilicone penetrant at the rate of 200-500 mL per 100 L of water
	Lantana ( <i>Lantana camara</i> )			For Lantana, apply when actively growing. DO NOT apply during periods of Summer drought stress. Use Holdfast Organosilicone penetrant at the rate of 200-500mL per 100 L of water.
	St Johns Wort ( <i>Hypericum perforatum</i> )			For St. John's Wort, apply when actively growing from Spring to Summer. Use Holdfast Organosilicone penetrant at the rate of 200-500 mL per 100 L of water.
	Sweet Briar ( <i>Rosa rubiginosa</i> )			For Sweet Briar, apply when in full leaf, prior to leaf fall. Use Holdfast Organosilicone penetrant at the rate of 200-500 mL per 100 L of water

\*\* LEGUMES IN THE PASTURE WILL BE DAMAGED

**NATIVE PASTURES, RIGHTS OF WAY, COMMERCIAL AND INDUSTRIAL AREAS, DOMESTIC AND PUBLIC SERVICE AREAS**

**CONTROL USING LOWER RATE OF GLYPHOSATE AND HIGHER RATE OF METSULFURON**

- WEED FORCE EVICT Bio 510 + RENOVATE BROADLEAF & WOODY WEED TANK MIX (metsulfuron-methyl 600g/kg)

(always refer to First Aid, Safety Directions, Resistant Weeds Warning and any other restrictions for Weed Force Renovate Herbicide)

**FOR ALL STATES**

**RESTRAINTS:**

**DO NOT** apply if rainfall is expected within 6 hours of spraying.

**DO NOT** store tank mixes with other products.

**DO NOT** store the prepared spray for more than 2 days.

**DO NOT** apply to plants wet with dew or rain or if droplets run-off when plants are disturbed.

**TABLE 12. Control using lower rate of glyphosate and higher rate of metsulfuron**

SITUATION	WEEDS CONTROLLED	STATES	RATE per 100 L HANDGUN OR KNAPSACK Always add a non-ionic surfactant unless otherwise specified	CRITICAL COMMENTS
				Results may be slow to appear and may not be visible for 3-6 weeks. Full brown-out may take 3-6 months.
Native Pastures **, Rights of Way, Commercial and Industrial Areas, Domestic and Public Service Areas	Apple Box ( <i>Angophora floribunda</i> )	NSW only	140 mL + 10 g Renovate + Holdfast Organosilicone penetrant (200 -500mL/100 L)	Spray plants up to 4 m high ensuring thorough coverage. Plants growing from large lignotubers may not be controlled
	Australian Blackthorn ( <i>Bursaria spinosa</i> )	NSW, Qld, Vic, Tas only	140 mL + 10 g Renovate	Spray to point of run off.
	Bitou Bush/Boneseed ( <i>Chysanthemoides monillifera</i> )	NSW, Qld, Vic, SA only		Ensure thorough coverage. Minimise contact with desirable species.
	Blackberry ( <i>Rubus spp.</i> )	All States		Ensure thorough coverage. Peripheral runners must also be covered to minimise regrowth. Apply between January and April when bushes are actively growing. TAS ONLY: Apply after petal fall. Do not apply to bushes bearing mature fruit.
Pastures planting interval – refer to Renovate herbicide label				

**TABLE 12 CONTINUED. Control using lower rate of glyphosate and higher rate of metsulfuron**

SITUATION	WEEDS CONTROLLED	STATES	RATE per 100 L HANDGUN OR KNAPSACK Always add a non-ionic surfactant unless otherwise specified	CRITICAL COMMENTS
Native Pastures **, Rights of Way, Commercial and Industrial Areas, Domestic and Public Service Areas	Messmate Stringybark ( <i>E. obliqua</i> ) Narrowleaf Peppermint ( <i>E. radiata</i> ) Blakely's Red Gum ( <i>E. blakelyi</i> ) Yellow Box ( <i>E. melliodora</i> )	NSW only	140 mL + 10 g Renovate + Holdfast Organosilicone penetrant (200 -500mL/100 L)	Results may be slow to appear and may not be visible for 3-6 weeks. Full brown-out may take 3-6 months.  Spray plants up to 4 m high. Trees growing from large lignotubers may not be controlled.
	Gorse ( <i>Ulex europaeus</i> )	NSW, Vic, Tas, SA only		Ensure thorough coverage and penetration of spray into the bush. Apply to plants up to 2 m tall.
Pastures planting interval – refer to Renovate herbicide label	Hawthorn ( <i>Crataegus leavigata</i> )	NSW only	140 mL + 10 g Renovate + Holdfast Organosilicone penetrant (200 -500mL/100 L)	Spray to point of run off.
	Lantana ( <i>Lantana camara</i> )	NSW, Qld only		Ensure thorough coverage and penetration of spray into the bush. Apply to plants up to 2 m tall.
	Privet ( <i>Ligustrum spp.</i> )	NSW, Qld only		Apply to bushes up to 3 m high. Ensure thorough coverage. If plants are not adequately covered plants may not be controlled.
	St John's Wort ( <i>Hypericum perforatum</i> )	NSW, Vic, SA, WA only		Spray to wet all foliage, but not to cause run-off.
	Sweet Briar ( <i>Rosa rubiginosa</i> )	NSW, VIC, Tas SA only		Do not apply after leaf fall has commenced or after the end of February. Spray foliage to point of run-off.
	Tree of Heaven ( <i>Allanther altissima</i> )	NSW only		

\*\* LEGUMES IN THE PASTURE WILL BE DAMAGED

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED**

## GENERAL INSTRUCTIONS

Evict Bio 510 Herbicide is a water soluble liquid herbicide. The product is non-selective and will control a wide range of emerged annual and perennial weeds. It provides no residual activity and is inactivated once it comes in contact with the soil.

Evict Bio 510 Herbicide is absorbed by plant leaves and green stems and is then translocated throughout the plant to the root system. The product inhibits a plant enzyme causing a breakdown in the metabolic pathway leading to death of the plant.

Visible effects of the product efficacy are gradual wilting, yellowing leading to complete plant browning.

For annual weeds, effects are usually apparent in 3-7 days and for perennial weeds up to 14-21 days.

The time taken for these effects to appear will vary depending on the speed of translocation; which will be dependent on climatic conditions such as temperature, moisture conditions etc. Best results will be obtained if plants are sprayed when they are actually growing and not under stress from factors such as disease, waterlogging, insect damage, drought stress etc. To ensure that the product is adequately absorbed by weeds it is recommended that spraying be delayed if rainfall is expected. Rain within 1 hour of application, which causes run-off, may require re-treatment. Rainfastness is reduced if weeds are not actively growing, under stress or conditions of low light intensity/darkness. The addition of a nonionic surfactant may improve rainfastness on annual grass weeds. Plants which are covered in dust or which are wet with dew should not be treated.

### Crop establishment

Where Evict Bio 510 Herbicide is used to control weeds prior to the establishment of a new crop or pasture it is important that the crop or pasture not be sown until a suitable seedbed is present.

Where a light cover of weeds has been sprayed, it may be possible to sow after 1 hour of daylight after application. Where a large amount of dead matter or trash is present the seedbed needs to be adequately prepared before crop or pasture sowing.

## MIXING AND APPLICATION

Evict Bio 510 Herbicide may be applied by boom spray, air, knapsack, handgun or wiper application.

### Boom Equipment

Spray volumes of 25-100 L water/ha are recommended with fan nozzles at a pressure of 240-280 kPa.

Boom height must be set to ensure double overlap of nozzle patterns at the top of the weed canopy.

### Wiper Equipment

Ropewick, canvas, carpet or felt applicators may be used to apply the product in the situation as per the Directions for Use table. Weeds should be at least 15cm above the crop and the wiper equipment should be operated at least 10cm above the crop. Best results are achieved at lower speeds (do not exceed 8 kph) and where two applications are made in opposite directions (double pass).

Where herbicide does not contact foliage (due to different levels of foliage) results may not be satisfactory and re-treatment may be required. Do not store a mixed solution for more than a couple of days.

Rate: Mix 700mL Evict Bio 510 Herbicide with 2 litres clean water.

## Aerial Equipment

Apply a minimum spray volume of 20 L/ha for Micronair and boom equipment. Nozzles and pressure settings should be selected to deliver a MEDIUM or MEDIUM-COARSE size droplet at the target (ASAE S572).

Aerial application is only recommended in pasture or fallow situations before establishment of a new crop or pasture, or in pre-harvest sorghum.

Use recommended rates on this label up to a maximum limit of 2.8 L/ha.

On sloping ground or hilly terrain the spraying height may vary, so it is recommended that spray volume may be increased to 30-80 L/ha and nozzles that produce a COARSE spray quality (ASAE S572).

High temperatures and/or low relative humidity cause excessive evaporation of spray droplets which may reduce results. When the temperature reaches 25°C, increase the water volume to at least 30 L/ha and the droplet size to at least COARSE droplet size (ASAE S572).

**DO NOT** apply Evict Bio 510 Herbicide by aircraft when temperature is above 30°C.

**DO NOT** use in intensive horticultural areas.

### Aerial Equipment (by Helicopter Only)

Helicopter applications must also comply with general Aerial Equipment instructions above.

Evict 510 Bio may be applied by helicopter for the control of weeds in forests, pasture or new forest plantings up to a maximum rate of 2.9 L/ha where specified by this label.

When applying this product by helicopter in combination with metsulfuron for the control of Blackberry in forestry and other specific situations, a rate of up to 5.6 L/ha of Evict Bio 510 may be applied.

Spray with a properly calibrated helicopter using the half overlap opposite pass technique and with a swath width set to minimise distribution variation.

Swath width will need to be reduced to avoid striping under light wind conditions and/or application to tall, dense targets.

Avoid spraying in still conditions and where conditions favour temperature inversions.

Turn off spray booms whilst passing over creeks and dams.

Thoroughly wash helicopter after each day of spraying to remove herbicide residues.

### Avoid Drift:

**DO NOT** use with spraying equipment or under meteorological conditions which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. Equipment settings which produce fine droplets (150 microns or less), winds over 12 km/h, temperature inversion conditions, still air and hot dry days all contribute to drift.

## **SURFACTANT**

The addition of surfactant may improve weed control where water rates are high or product rates low. Suggested surfactant rates are 200mL/100L of 1000g/L non-ionic surfactant or 250-500mL of 700g/L surfactant (eg Sidekick Wetting Agent & Spray Conditioner. Do not add spraying oils, agricultural chemicals or other materials except as directed on the label.

Woody & perennial weeds:

When spraying woody and perennial weeds use Holdfast Organosilicone penetrant at the rate of 200-500mL/100L.

Add Sidekick Wetting Agent & Spray Conditioner at RATE: 300-500mL/100L spray solution

## **MIXING**

When the product is to be mixed with water it is important that clean water is used. Dirty water or hard water containing calcium salts may reduce the product's effectiveness.

The following procedure should be followed:

1. Ensure spray tank is clean and that the previous chemicals used are washed from the tank.
2. Half fill the tank with clean water and the required amount of Evict Bio 510 Herbicide.
3. Add the rest of the water.
4. Add surfactant last.

## **COMPATABILITY**

Evict Bio 510 Herbicide may be mixed with a wide variety of products to broaden the spectrum of weed control, and to add residual control. Refer to the 'Directions for Use' Section for detailed information on tank mix situations.

## **Additives:**

Ammonium sulphate: may be used as an adjuvant to alleviate the adverse effects of high levels of calcium, magnesium and bicarbonate ions in water. Rate – 2 L/ha spray solution.

## **Herbicides:**

### **Atrazine/Simazine**

Evict Bio 510 Herbicide may be mixed with Atrazine Flowable or Simazine Flowable for knockdown and residual weed control.

Addition of ammonium sulphate at 2% w/v (2 kg/100 Litres of spray solution) is recommended to avoid antagonism.

### **Weed Stop Herbicide:**

Evict Bio 510 Herbicide may be mixed with Weed Stop herbicide for knockdown and residual weed control. Refer to the Weed Stop label for details.

### **Ground Keeper Herbicide (sulfometuron):**

Evict Bio 510 Herbicide may be mixed with Ground Keeper Herbicide for knockdown and residual weed control. Refer to the Ground Keeper label for details.

### **Xtract Herbicide (clopyralid):**

Evict Bio 510 Herbicide may be mixed with Xtract Herbicide for knockdown and residual weed control of fleabane and other weeds. Refer to the Xtract label for details.

### **Dicamba:**

Evict Bio 510 Herbicide and dicamba may be tank mixed for more effective control of Sorrel, Sub. clover and medics.

### **2,4-D:**

Evict Bio 510 Herbicide and 2,4-D ester or 2,4-D isopropylamine may be tank mixed for improved control of broadleaf weeds.

### **Chlorsulfuron:**

Evict Bio 510 Herbicide and chlorsulfuron tank mix will provide knockdown and residual control in fallow and crop. Observe plantback restrictions for chlorsulfuron.

### **Renovate herbicide(Metsulfuron-methyl):**

Evict Bio 510 Herbicide and metsulfuron tank mix will provide knockdown weed control in fallows, prior to planting certain winter cereals and forestry site preparation. Follow all instructions on the metsulfuron label. Observe plantback restrictions for metsulfuron.

### **Oxyfluorfen:**

The addition of oxyfluorfen at 75 mL/ha to recommended rates of Evict Bio 510 Herbicide used prior to planting wheat or barley will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity.

### **Insecticides:**

Chlorpyrifos, dimethoate, fenitrothion, omethoate and lambda-cyhalothrin. Other insecticides have not been tested.

## COMPATABILITY AGENT – Liquid ammonium sulphate

Rate: 200mL/100L spray solution. When mixing with oxyfluorfen, add to improve the compatibility in cold water (less than 15°C). Liquid ammonium sulphate must be pre-mixed with oxyfluorfen before adding to the spray tank. Refer to Liquid ammonium sulphate label for full directions.

### For tank mixing, the following procedure should be undertaken:

1. Half fill tank and start agitation.
2. Add ammonium sulphate.
3. Add companion product.
4. Add Evict Bio 510 Herbicide and the rest of the water.
5. Add surfactant and maintain agitation while spraying.

## EQUIPMENT MAINTENANCE AND USAGE

Evict Bio 510 Herbicide should only be stored, mixed or applied in plastic, or plastic-lined, stainless steel, aluminium, copper, brass or fibreglass containers. The product and spray solutions react with galvanised steel and unlined steel tanks and containers to form hydrogen gas, which may form a highly combustible gas mixture. This gas could cause an explosion if ignited by an open flame. All application equipment, including tanks, nozzles, hoses, aircraft and aircraft landing gear, should be thoroughly washed after use to prevent corrosion.

## RESISTANT WEEDS WARNING

<b>GROUP</b>	<b>9</b>	<b>HERBICIDE</b>
--------------	----------	------------------

Evict Bio 510 Herbicide is a member of the Glycine group of herbicides. Evict Bio 510 Herbicide has the inhibitor of EPSP syntheses mode of action. For weed resistance management Evict Bio 510 Herbicide is a Group 9 Herbicide.

Some naturally occurring weed biotypes resistant to Evict Bio 510 Herbicide and other Group 9 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Evict Bio 510 Herbicide or other Group 9 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Weed Force Pty Ltd accepts no liability for any losses that may result from the failure of Evict Bio 510 Herbicide to control resistant weeds.

## PROTECTION OF CROP, NATIVE AND OTHER NON-TARGET PLANTS

This product is non-selective and may severely injure or kill desirable plants should the product come into contact with foliage, green stems or fruit of such plants. DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures. DO NOT allow spray to contact any part of desirable plants. DO NOT use prior to transplanting tomato seedlings.

## PROTECTION OF LIVESTOCK

There is no withholding period for this product, but removal of stock may be necessary to achieve efficacy. It is recommended that stock be removed from the area to be treated 1 day before treatment of annual weeds and 7 days for perennial weeds. Certain plants (eg soursob, variegated thistle) may be naturally toxic to stock. Where known toxic plants are present, do not allow stock to graze until complete browning of treated plants has occurred.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, streams, waterways or rivers with the chemical or used containers. When controlling weeds near water, refer to label directions to minimise the entry of spray into the water.

## PRECAUTION

DO NOT store, mix or apply the product or spray solutions in unlined steel or galvanised containers as a highly flammable gas may form. Use stainless steel, brass, copper, aluminium, plastic or plastic lined, fibreglass containers or spray tanks.

## STORAGE AND DISPOSAL

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

Product will irritate eyes and skin. Avoid contact with eyes and skin. When preparing the product for use, wear elbow length PVC gloves and face shield or goggles. When using controlled droplet applicator, wear protective waterproof clothing and impervious footwear.

After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## SAFETY DATA SHEET (SDS)

The SDS contains important information relating to health and safety.

This product should be used in conjunction with information on the SDS and users are advised to read the SDS before using this product. The most up to date SDS can be obtained from [www.weedforce.com.au](http://www.weedforce.com.au)



**CONDITIONS OF SALE**

To the extent permitted by law, all conditions and warranties and statutory or other rights or action which buyer or any other user may have against Weed Force Pty Ltd ("Weed Force") or Seller are hereby excluded.

Weed Force hereby gives Notice to Buyer and other users that it will not accept responsibility for any direct, indirect or consequential loss however arising (including negligence) from the sale, supply, use or application of this product or reliance on product information or advice provided by Weed Force or on its behalf. No warranty expressed or implied is given by Weed Force regarding its suitability or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not. Weed Force's liability shall, in all circumstances, be limited to replacement of the product or a refund of the purchase price paid therefor.



WEEDFORCE Pty Ltd ACN 602 207 152   
2072 Logan Rd Mt Gravatt QLD 4122  
P: 0488 997 923 [www.weedforce.com.au](http://www.weedforce.com.au)

® Trademark of Weed Force Pty Ltd  
© Weed Force Pty Ltd 2017

APVMA Approval No:  
86155/118321